

For adults living with narcolepsy...
Now FDA approved for cataplexy

Wakix[®]
pitolisant tablets

WAKIX is a prescription medicine approved for the treatment of excessive daytime sleepiness (EDS) or cataplexy in adults with narcolepsy

Why WAKIX?



WAKIX is a **first-of-its-kind medication** that increases histamine levels in the brain.

The way that WAKIX works to treat **EDS** or **cataplexy** in adults with narcolepsy is not fully understood.



WAKIX is the **only** FDA-approved treatment for people with narcolepsy that is **not a controlled substance**.



WAKIX is the **only once-daily tablet** that is **FDA approved to treat EDS or cataplexy in adults with narcolepsy**.



WAKIX is **not a stimulant**.

Important Safety Information

Do not take WAKIX if you are allergic to pitolisant or any ingredient in WAKIX, or if you have severe liver disease.

Tell your healthcare provider about all your medical conditions, including if you have heart rhythm irregularities, were born with a heart condition, or the levels of electrolytes in your blood are too high or too low. WAKIX has an effect on the electrical activity of the heart known as QT/QTc prolongation. Medicines with this effect can lead to disturbances in heart rhythm, which are more likely in patients with risk factors such as certain heart conditions, or when taken in combination with other medicines that affect QT. Tell your healthcare provider about all the other medicines you take.

The risk of QT prolongation may be greater in patients with liver or kidney disease. WAKIX is not recommended in patients with end-stage kidney disease.

The most common side effects seen with WAKIX were insomnia, nausea, and anxiety. Other side effects included headache, upper respiratory infection, musculoskeletal pain, heart rate increased, and decreased appetite. These are not all the possible side effects of WAKIX. Tell your healthcare provider about any side effect that bothers you or that does not go away.

Tell your healthcare provider about all the medicines you take or plan to take, including prescription and over-the-counter medicines. Some medicines can increase the amount of WAKIX that gets into your blood and some medicines can decrease the amount of WAKIX that gets into your blood. The dosage of WAKIX may need to be adjusted if you are taking these medicines.

WAKIX can also decrease the effectiveness of some medicines, including hormonal birth control methods. You should use an alternative non-hormonal birth control method during treatment with WAKIX and for at least 21 days after discontinuation of treatment.

Tell your healthcare provider if you are pregnant or planning to become pregnant. There is a pregnancy exposure registry that monitors pregnancy outcomes in women who are exposed to WAKIX during pregnancy. You are encouraged to enroll in the WAKIX pregnancy registry if you become pregnant while taking WAKIX. To enroll or obtain information from the registry, call 1-800-833-7460.

The safety and effectiveness of WAKIX have not been established in patients less than 18 years of age.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088. You can also report negative side effects to Harmony Biosciences at 1-800-833-7460.

Please see accompanying Full Prescribing Information.



**Ask your healthcare provider if WAKIX might be right for you.
Visit WAKIX.com for more information.**



WAKIX is a registered trademark of Bioprojet Pharma.
Harmony Biosciences name and logo are registered trademarks.
© 2020 Harmony Biosciences. All rights reserved.
US-WAK-2000317/Oct 2020